

# Singularities in axially symmetric solutions of Einstein-Yang Mills and related theories

Ludger Hannibal

*Fachbereich Physik, Carl von Ossietzky Universität Oldenburg, D-26111*

*Oldenburg, Germany*

(March 8th, 1999)

We show that the solutions of  $SU(2)$  Yang-Mills-dilation and Einstein-Yang-Mills-dilaton theories described in a sequence of papers by Kleihaus and Kunz are not regular in the gauge field part.

Since the pioneering papers by Bogomol'nyi [1] and Witten [2] on exact solutions of classical Yang-Mills theory many studies in this field have been carried out. In recent time the gravitational self-interaction of solitons has been of interest [3]. Not only spherically symmetric solutions of theories including Yang-Mills fields, with or without gravitation, Higgs, or dilaton fields, have been constructed, but also solutions with axial symmetry [4]. These latter type of solitons was studied by Kleihaus and Kunz in the context of Yang-Mills-dilaton theory [5] and Einstein-Yang-Mills-dilaton theory [6–9], mainly by numerical construction. The relevance of solutions containing black holes is that they should provide counterexamples to the no-hair conjecture for such theories. This conjecture is the analog to the no-hair theorems in Einstein-Maxwell and related theories [10].

In this communication we would like to point out that the gauge field part of the solutions described in Refs. [5–9] are not regular on the symmetry axis, in contradiction to what is claimed. Thus the solutions are similar to singular monopole solutions and thus have to be interpreted carefully [11–13]. We explicitly derive conditions for the regularity, and show that a recent attempt by Kleihaus [15] to relate the non-regular solution to regular ones via a (non-regular) gauge transformations like it is done in the case of monopoles falls short of its aim, so that one must ask whether the solutions described exist in any well-defined way.

Non-regular gauge fields are encountered within Dirac's theory of monopoles. In general, solutions of Maxwell's equations for the electromagnetic four-potential must be twice continuous differentiable in order to be acceptable from the mathematical point of view. From the Aharonov-Bohm effect we know that continuity of the potential is essential. For the derivation of the field equations from an action one uses partial integration, thus the solutions must be differentiable. In general, regularity means that all solutions are at least twice differentiable in the sense of distributions, allowing for delta-like sources. Vacuum solution will in general be locally analytic. In Dirac's theory of monopoles there ex-

ists no globally defined regular gauge field, only the electromagnetic field tensor is regular. One can find a gauge transformation in the vicinity of the singularity on the symmetry axis that gives a locally regular gauge field, and the compatibility condition in the overlap region leads to the quantization of the monopole [11,12]. In the case of non-abelian gauge theories the situation is more complicated, and for overlapping solutions connected by gauge transformations compatibility conditions are required [13]. Another problem is that the gauge potential explicitly enter in the field equations for the gauge field tensor, because in contrast to the abelian case the gauge field tensor is only gauge covariant, not invariant. Thus, at singular points of the gauge potential the field equations are not well defined, even if the field tensor is. In the case of spherically symmetric solutions there will in general be only a singularity at the origin, the coordinate singularity on the  $z$ -axis plays no role due to the spherical symmetry. Witten [2] explicitly showed how such a singularity can be removed by a gauge transformation.

In the case of axially symmetric solutions the situation becomes more complicated, since a coordinate singularity at  $\theta = 0, \pi$  is introduced, but cannot be removed by a symmetry argument. Together with axial symmetry Rebbi and Rossi [4] introduced a winding number  $n$  into their ansatz for the gauge field, using unit vectors in the Lie algebra  $su(2)$  of  $SU(2)$  that correspond to cylindrical coordinates  $(\rho, \phi, z)$ . Writing these as

$$\begin{aligned} u_1 &= \tau_\rho^n = \tau_1 \cos n\phi + \tau_2 \sin n\phi \\ u_2 &= \tau_\phi^n = -\tau_1 \sin n\phi + \tau_2 \cos n\phi \\ u_3 &= \tau_z^n = \tau_3 \end{aligned} \quad (1)$$

where  $\tau_i, i = 1, 2, 3$  are the Pauli matrices, the gauge field attains the form

$$A_\mu = u_i W_\mu^i \quad (2)$$

with twelve independent gauge fields  $W_\mu^i, i = 1, 2, 3, \mu = 0, 1, 2, 3$ . In spherical coordinates  $(r, \theta, \phi)$  the unit vector of  $su(2)$  will be

$$\begin{aligned} \tau_r^n &= \tau_1 \sin \theta \cos n\phi + \tau_2 \sin \theta \sin n\phi + \tau_3 \cos \theta, \\ \tau_\theta^n &= \tau_1 \cos \theta \cos n\phi + \tau_2 \cos \theta \sin n\phi - \tau_3 \sin \theta, \\ \tau_\phi^n &= -\tau_1 \sin n\phi + \tau_2 \cos n\phi \end{aligned} \quad (3)$$

The gauge potentials  $A_\mu$  are not regular unless the functions  $W_\mu^i$  behave appropriately on the  $z$ -axis, since the functions  $\sin n\phi, \cos n\phi$ , when considered as functions on  $S^2$ , are neither continuous nor differentiable at the poles

$\theta = 0, \pi$ . The functions  $\sin n\phi$  and  $\cos n\phi$  can be expanded into polynomials of degree  $|n|$  in the variables  $\sin \phi = y/\rho$  and  $\cos \phi = x/\rho$ , with  $\rho = \sqrt{x^2 + y^2}$ ; all terms of these polynomials have denominators that are even powers of  $\rho$  for  $n$  even, and odd powers of  $\rho$  for  $n$  odd. Hence  $\rho^{|n|} \sin n\phi$  and  $\rho^{|n|} \cos n\phi$  are regular functions except for  $r = 0$ , and may further be multiplied by functions of the variables  $\rho^2$  and  $z$  without losing their regularity. This was explicitly taken into account by Rebbi and Rossi [4]. In spherical coordinates functions of the type

$$f(r, \theta^2) \sin^{|n|} \theta \sin n\phi + g(r, \theta^2) \sin^{|n|} \theta \cos n\phi \quad (4)$$

with regular functions  $f$  and  $g$  are regular along the axis  $\theta = 0$ , possibly except at  $r = 0$ . The well defined  $\theta$ -parity of these functions is easily understood. On any great circle through the pole at  $\theta = 0$ , with  $\phi = \phi_0$  to the right of the pole,  $\phi = \phi_0 \pm \pi$  to the left, and coordinate

$$\alpha = \begin{cases} \theta & \text{for } \alpha > 0, \phi = \phi_0 \\ -\theta & \text{for } \alpha < 0, \phi = \phi_0 \pm \pi \end{cases} \quad (5)$$

we have for any function  $F(\theta)$  (defined only for  $\theta > 0$ )

$$\begin{aligned} \lim_{\alpha \nearrow 0} \frac{\partial^k}{\partial \alpha^k} F(\theta = -\alpha) \sin n(\phi_0 \pm \pi) \\ = (-1)^{n+k} \lim_{\alpha \searrow 0} \frac{\partial^k}{\partial \alpha^k} F(\theta = \alpha) \sin n\phi_0. \end{aligned} \quad (6)$$

Therefore it is necessary that  $F(\theta)$  is an even function in  $\theta$  for  $n$  even, an odd function for  $n$  odd in order that  $F(\theta) \sin n\phi$  can be continuous ( $k = 0$ ) or that the derivatives exist ( $k = 1, 2, \dots$ ). Together with the multiplication by the factor  $\sin^{|n|} \theta$  we obtain (4) as a sufficient form for functions to be analytical at  $\theta = 0$ .

Kleihaus and Kunz use spherical coordinates  $(r, \theta, \phi)$ . Their ansatz for the  $SU(2)$  gauge field is [8]:

$$A_\mu dx^\mu = \frac{1}{2gr} \left\{ \begin{aligned} &\tau_\phi^n [H_1 dr + (1 - H_2) r d\theta] \\ &- n [\tau_r^n H_3 + \tau_\theta^n (1 - H_4)] r \sin \theta d\phi \end{aligned} \right\} \quad (7)$$

This gauge potential is singular at  $x = y = 0$  unless appropriate conditions are imposed on the functions  $H_i$ ,  $i = 1, 2, 3, 4$  that depend only on  $r$  and  $\theta$ . In order to analyze this ansatz for regularity, we decompose the gauge potential into its Cartesian components,

$$A_i dx^i = \frac{1}{2gr} \sum_{j=1}^3 \tau^j \left( w_1^j dx + w_2^j dy + w_3^j dz \right), \quad (8)$$

with the nine functions  $w_i^j$  given by

$$\begin{aligned} w_1^1 &= -F_1 \sin n\phi \cos \phi + nF_3 \cos n\phi \sin \phi \\ w_1^2 &= F_1 \cos n\phi \cos \phi + nF_3 \sin n\phi \sin \phi \\ w_1^3 &= nF_4 \sin \phi \end{aligned}$$

$$\begin{aligned} w_2^1 &= -F_1 \sin n\phi \sin \phi - nF_3 \cos n\phi \cos \phi \\ w_2^2 &= F_1 \cos n\phi \sin \phi - nF_3 \sin n\phi \cos \phi \\ w_2^3 &= -nF_4 \cos \phi \\ w_3^1 &= -F_2 \sin n\phi \\ w_3^2 &= F_2 \cos n\phi \\ w_3^3 &= 0. \end{aligned} \quad (9)$$

The functions  $F_1$  to  $F_4$  are defined by

$$\begin{aligned} F_1 &= H_1 \sin \theta + (1 - H_2) \cos \theta \\ F_2 &= H_1 \cos \theta - (1 - H_2) \sin \theta \\ F_3 &= H_3 \sin \theta + (1 - H_4) \cos \theta \\ F_4 &= H_3 \cos \theta - (1 - H_4) \sin \theta. \end{aligned} \quad (10)$$

We see that a sufficient condition that the gauge potential can be analytically extended to  $\theta = 0$  for  $r > 0$  is

$$F_1(r, \theta) = \tilde{F}_1(r, \theta^2) \sin^{|n|+1} \theta \quad (11)$$

$$F_2(r, \theta) = \tilde{F}_2(r, \theta^2) \sin^{|n|} \theta \quad (12)$$

$$F_3(r, \theta) = \tilde{F}_3(r, \theta^2) \sin^{|n|+1} \theta \quad (13)$$

$$F_4(r, \theta) = \tilde{F}_4(r, \theta^2) \sin \theta \quad (14)$$

with analytic functions  $\tilde{F}_1$  to  $\tilde{F}_4$ . If regularity is sought only up to a certain number of derivatives, then accordingly less constraints on the odd orders in  $\theta$  are needed. Now a more subtle point is that these conditions guarantee regularity, but are not necessary. In the special case  $n = 1$  with spherical symmetry and  $H_1 = H_3 = 0, H_2 = H_4 = w(r)$  we see that  $F_1$  and  $F_3$  do not respect (11) and (13), respectively. The reason is that in the case  $F_1 = F_3$  the relation  $\cos^2 \phi + \sin^2 \phi = 1$  leads to a regular function where in general  $F_1 \cos^2 \phi + F_3 \sin^2 \phi$  would not have been regular. It is not difficult to see that for arbitrary  $n$  this will happen if

$$F_1 = nF_3 \quad (15)$$

so that we can do with two powers less of  $\sin \theta$  in (11) and (13) if the functions agree in the lowest order. Hence an ansatz that is analytical and also respects all symmetry requirements at  $\theta = 0, \pi/2, \pi$  is given by

$$F_1(r, \theta) = \left[ nf(r) + \tilde{F}_1(r, \sin^2 \theta) \sin^2 \theta \right] \cos \theta \sin^{|n|-1} \theta \quad (16)$$

$$F_2(r, \theta) = \tilde{F}_2(r, \sin^2 \theta) \sin^{|n|} \theta \quad (17)$$

$$F_3(r, \theta) = \left[ f(r) + \tilde{F}_3(r, \sin^2 \theta) \sin^2 \theta \right] \cos \theta \sin^{|n|-1} \theta \quad (18)$$

$$F_4(r, \theta) = \tilde{F}_4(r, \sin^2 \theta) \sin \theta \quad (19)$$

For the function  $H_i$  this implies

$$\begin{aligned} H_1 &= -F_1 \sin \theta + F_2 \cos \theta \\ &= \left[ -nf(r) - \tilde{F}_1 \sin^2 \theta + \tilde{F}_2 \right] \cos \theta \sin^{|n|} \theta \\ (1 - H_2) &= F_1 \cos \theta + F_2 \sin \theta \end{aligned}$$

$$\begin{aligned}
&= \left[ \left( n f(r) + \tilde{F}_1 \sin^2 \theta \right) \cos^2 \theta + \tilde{F}_2 \sin^2 \theta \right] \sin^{|n|-1} \theta \\
H_3 &= -F_3 \sin \theta + F_4 \cos \theta \\
&= \left[ \left( -f(r) - \tilde{F}_3 \sin^2 \theta \right) \sin^{|n|-1} \theta + \tilde{F}_4 \right] \sin \theta \cos \theta \\
(1 - H_4) &= F_3 \cos \theta + F_4 \sin \theta \\
&= \left[ f(r) + \tilde{F}_3 \sin^2 \theta \right] \cos^2 \theta \sin^{|n|-1} \theta + \tilde{F}_4 \sin^2 \theta. \quad (20)
\end{aligned}$$

This especially implies that  $H_1$  and  $1 - H_2$  will always have a well defined  $\theta$ -parity, whereas for  $H_3$  and  $1 - H_4$  this is the case only for odd  $n$ . At the point  $r = 0$  the gauge potential is analytical, if moreover the functions  $\tilde{F}_i$  have the form

$$\begin{aligned}
\tilde{F}_i(r, \sin^2 \theta) &= r^2 \check{F}_i(r^2, \sin^2 \theta) \text{ for } i = 1, 3 \\
\tilde{F}_i(r, \sin^2 \theta) &= r \check{F}_i(r^2, \sin^2 \theta) \text{ for } i = 2, 4 \quad (21)
\end{aligned}$$

with analytical functions  $\check{F}_i$ . The solutions constructed by Kleihaus and Kunz [5–9] do not have the regular form (20) [14].

Kleihaus most recently showed [15] that for  $n = 2, 3, 4$  a gauge transformation of the form

$$U(\vec{r}) = \exp(i \frac{\Gamma}{2} \tau_\varphi^n) \quad (22)$$

leads to a new gauge field

$$\begin{aligned}
\hat{A} &= U A U^\dagger + i dU U^\dagger \\
&= \frac{1}{2gr} \left\{ \begin{array}{l} \tau_\varphi^n \left[ \hat{H}_1 dr + (1 - \hat{H}_2) r d\theta \right] \\ -n \left[ \hat{F}_3 \tau_\rho^n + \hat{F}_4 \tau_3 \right] r \sin \theta d\phi \end{array} \right\} \quad (23)
\end{aligned}$$

that satisfies the following conditions:

$$\begin{aligned}
\hat{H}_1 &\sim \rho^{|n|} + O(\rho^{|n|+1}) \\
1 - \hat{H}_2 &\sim \rho^{|n|+1} + O(\rho^{|n|+2}) \\
\hat{F}_3 &= \left[ \frac{\rho}{r} \hat{H}_3 + \frac{z}{r} (1 - \hat{H}_4) \right] \sim \rho^{|n|+1} + O(\rho^{|n|+2}) \\
\hat{F}_4 &= \left[ \frac{z}{r} \hat{H}_3 - \frac{\rho}{r} (1 - \hat{H}_4) \right] \sim \rho + O(\rho^2) \quad (24)
\end{aligned}$$

We see that this transformation resulted in a continuous gauge potential [16], but since the conditions on the  $\theta$ -parity have not been considered, the potentials are still possibly not differentiable at  $\theta = 0$ . One must at least show that the next to leading order vanishes, then the potentials will be twice differentiable at least. Especially for even  $n$  the  $\theta$ -parity of the functions  $H_1$  and  $H_2$ , which is necessary in view of (10) and (16) and (17), is not respected in the expansions given. Thus the desired aim, namely to show that one can obtain all gauge invariant quantities also from a regular gauge potential, that exists at least locally, has been failed. Moreover, where expansions of the gauge potential functions  $H_i$  have been carried without respecting the necessary  $\theta$ -parity, we must indeed ask whether regular solutions with axial symmetry

exist at all. A proof of existence is certainly needed, especially since more boundary conditions must be satisfied than can be given to the numerical routines for solving the nonlinear partial differential equations for these functions. Therefore, at least so far as the  $SU(2)$  Einstein-Yang-Mills(-dilaton) theory has been studied by Kleihaus and Kunz, the numerical solutions give insufficient indication for the existence of corresponding exact solutions.

Finally, a remark about the interpretation of the theory appears to be indicated. Provided the existence of regular solutions has been established, their discussion should not be based on gauge-dependent properties of some singular potential. We have seen that the function  $f$ , which is the only relevant one in the spherically symmetric case, has no physical meaning in the axially symmetric case [16]. For  $n > 1$  the functions  $H_2$  and  $H_4$  must satisfy  $H_2 = H_4 = 1$  on the symmetry axis for any regular gauge potential, which means that the node number of these functions introduced in the discussion are artifacts of the singular gauge. Hence the classification of solutions that are not equivalent under gauge transformations must be based on some other aspect.

- 
- [1] E. B. Bogomol'nyi, Sov. J. Nucl. Phys. **24** 861 (1976)
  - [2] E. Witten, Phys. Rev. Lett. **38** (1977) 121
  - [3] M. S. Volkov and D. V. Gal'tsov "Gravitationg Non-Abelian Solitons and Black Holes with Yang-Mills Fields" preprint hep-th/9810070
  - [4] C. Rebbi and P. Rossi, Phys. Rev. D **22**, 2010 (1980)
  - [5] B. Kleihaus and J. Kunz, Phys. Lett. B **392**, 135 (1997)
  - [6] B. Kleihaus and J. Kunz, Phys. Rev. Lett. **78**, 2527 (1997)
  - [7] B. Kleihaus and J. Kunz, Phys. Rev. Lett. **79**, 1595 (1997)
  - [8] B. Kleihaus and J. Kunz, Phys. Rev. D **57**, 834 (1998)
  - [9] B. Kleihaus and J. Kunz, Phys. Rev. D **57**, 6138 (1998)
  - [10] see: M. Heusler, *Black Hole Uniqueness Theorems* (Cambridge, Cambridge University Press, 1996) for an overview
  - [11] G. Wentzel, Progr. Theor. Phys.(Kyoto) Suppl. **37** and **38** (1966) 163.
  - [12] J. Arafune, P. G. O. Freund and C. J. Goebel, J. Math. Phys. **16** (1975) 433.
  - [13] T. T. Wu and C. N. Yang, Phys. Rev. D **12** 3857 (1975).
  - [14] As was pointed out by the author on occasion of a talk given by B. Keihaus in Oldenburg in 1998.
  - [15] B. Kleihaus "On the regularity of static axially symmetric solutions in  $SU(2)$  Yang-Mills-dilaton theory" preprint hep-th/9901096, to be published in Phys. Rev. D
  - [16] The function  $f$  can be removed by a gauge transformation with  $\Gamma = -f(r) \sin^{|n|} \theta$ .